

Asymmetries in Powerful Extragalactic Radio Sources

Gopal-Krishna

*National Centre for Radio Astrophysics, Tata Institute of Fundamental Research, Pune University Campus,
Post Bag No. 3, Pune 411 007, India*

Paul J. Wiita

Department of Physics & Astronomy, Georgia State University, P.O. Box 4106, Atlanta GA 30302-4106, USA

My own suspicion is that the universe is not only queerer than we suppose, but queerer than we can suppose — J.B.S. Haldane

Overview

This review aims to provide the background and current perspective of some of the striking asymmetries exhibited by extragalactic double-lobed radio sources, paying special attention to the reported interrelationships among some of them. The asymmetries addressed here are related not only to the jets, and we particularly aim to highlight some less discussed aspects pertaining to the radio lobes and hot-spots for which any relativistic beaming effects are milder than those imposed on the jets. Various attributes of these features, such as their geometry, intensity, spectrum, and polarization provide important underpinnings to models of radio galaxies and quasars, as well as to theories that attempt to unify them. These constraints can be further sharpened by exploiting the more recently discovered asymmetry properties of the extended optical emission-line regions (EELR) associated with powerful radio sources, including the giant halos of Lyman- α emission associated with them. We suggest that highly extended disks of dusty gaseous material, which could also be parts of cosmic filaments, around the host galaxies of radio sources may hold the key to understanding some important types of asymmetries, namely, the correlated radio-optical asymmetry and perhaps also the depolarization asymmetry of the twin radio lobes.

We also discuss the nature and possible implications of the extreme type of asymmetry witnessed in the enigmatic quasar 3C273, whose *entire* radio emission is observed on just one side of the nucleus. In this context, theoretical scenarios for the production of one-sided jets are also briefly recounted. Taking a clue from the observed one-sidedness of the extended radio jets even in ‘weak-headed-quasars’, we also propose a scenario for the late evolutionary stage of double radio sources. We end by providing a list of key questions relating to radio source asymmetries which still require final answers.

1. Background and General Perspective

Although a bifurcated ejection of energy from radio-loud active galactic nuclei (AGN) in two oppositely directed channels had come to be appreciated already by the 1960s from the observed similarity and roughly symmetric disposition of the twin radio lobes about the parent galaxy, many different forms of departures from this symmetry have been discovered during the subsequent three decades. Investigations of the systematics and interrelations between the various manifestations of *asymmetry*

have played a pivotal role in the development of the present physical framework for extragalactic double radio sources, as discussed in the next sections. With the widespread application of CCD detectors during the last decade, the search for structural asymmetries could be successfully extended to much shorter wavebands, namely, the ultra-violet, optical, near-infrared, and even X-rays. The images obtained through the different electromagnetic windows have greatly enlarged our vision of the asymmetries, firstly by revealing new types of asymmetries and, furthermore, by exposing trends that signify important correlations among different manifestations of the asymmetries. This encouraging backdrop has provided a strong impetus for the studies that could sharpen the theories of double radio sources by confronting them with the rapidly widening domain of the observational findings in this area.

The earliest inkling of structural asymmetry came from the conspicuous one-sidedness of the optical jet already noted by Curtis (1918) in the nearby galaxy M87, which was later demonstrated to be a synchrotron source (Baade 1956), and whose apparent asymmetry has since been clearly demonstrated in the radio and some other wavebands as well. The double-lobed radio structure of such radio galaxies was first discovered, unexpectedly, from radio interferometric observations of the powerful radio galaxy Cygnus A (Jennison & Das Gupta 1953), which led to the concept of (non-spherical) symmetry of radio sources. Thus, in a sense, the concern over an asymmetric characteristic actually predates the discussions about the symmetric attributes of external galaxies (e.g., the double-lobed radio morphology).

Since asymmetry of jets is found to be a ubiquitous feature of quasars, the question of ‘hiding’ the jet feeding the other radio lobe led to the ‘relativistic beaming’ paradigm for the jets (e.g., Rees et al. 1982 and references therein), culminating in the development of the unified scheme, according to which quasars and radio galaxies are the same in kind and differ merely in orientation of the principal axis from our line-of-sight. While the relativistic jets are highly sensitive to orientation effects, the apparent properties such as the relative brightness and the locations of the slower moving components, including the radio lobes and hot-spots, can also be measurably influenced by orientation effects, on account of light-travel-time differences and Doppler boosting. Thus, the asymmetries have been effectively used to infer the dynamics, orientation, intensity evolution and ages of such components, laying the boundary conditions for theoretical models of them. Additionally, the red-shift dependences of the structural asymmetry and distortions of double radio sources have been investigated, in order to probe the cosmological evolution of the radio source properties and their environment.

The potential of the radio source asymmetry received a major boost in 1988, following the discovery of the depolarization asymmetry of the radio lobe pair of quasars (§3). This so called Laing–Garrington effect is generally believed to be also an orientation effect, and has been instrumental in clinching the long-standing debate in favor of the postulated relativistic bulk motion in the *kiloparsec-scale* radio jets of very powerful radio sources. At the moment, the status and understanding of the spectral index asymmetry, first reported in 1991 (§4), is rather confusing. From the available results based on rather small samples, it is still unclear to what extent the asymmetry must be intrinsic to the sources.

A new dimension was added to such studies with the discovery of the tightly correlated nature of the structural asymmetries revealed by the nonthermal radio and the thermal optical line emission. Two radically different perspectives of this phenomenon are discussed in §5. Another recently discovered optical asymmetry in high- z radio sources is related to the giant halos of Ly- α emission associated with the radio lobes, which are probably excited by the UV photons from the AGN. The apparent morphological asymmetries of the Ly- α halos (together with their observed kinematical profiles) can provide a powerful means of inferring the orientations of the radio sources, the motions of the host galaxies, and, in particular, of estimating the physical properties of the dusty superdisks

which appear to be commonly associated with high- z radio galaxies (RGs) (§§5 & 6). Aside from being the tracers of dust in the early universe, these superdisks could be instrumental in causing several prominent types of asymmetry.

Observations of asymmetries have proved to be a strong catalyst for theoretical studies and numerical simulations concerned with the production and propagation of the AGN beams, including models for ejecting one-sided (alternating) jets. Although not favored currently, such possibilities must still be kept in sight, and in §7 we provide some justification for this cautious approach. Despite our brief consideration of intrinsic jet asymmetries, we stress that in this review we are mainly concerned with the asymmetry of mildly beamed or unbeamed components (the hot-spots and lobes). Thus we do not discuss here the huge number of papers focussed on jet asymmetry.

2. The radio lobe-length and flux density asymmetries in FR II sources

2.1 Historical perspective

Although the term “classical double radio source” stems from the roughly symmetric appearance of the two radio lobes about the parent active galaxy, it is the prevailing degree of the observed asymmetry that holds the key to constraining many aspects of their physical models and their unification theories. The potential of such studies has been appreciated and exploited throughout much of the last four decades. Thus, while developing one of the earliest (kinematical) models of double radio sources, Ryle & Longair (1967) pointed out that light-travel-time effects and Doppler boosting due to relativistic velocities of the radio lobes can cause an apparent asymmetry of the lengths and flux densities of the twin lobes, if the source axis happens to be oriented away from the sky plane. De Young & Axford (1967) gave an early account of the environmental impact in terms of ram pressure confinement of the lobes. Within the framework of a simple kinematical model, the apparent lobe-length ratio (the asymmetry parameter, $Q \equiv D_{\text{approach}}/D_{\text{recede}}$) and the apparent flux ratio ($R \equiv F_{\text{approach}}/F_{\text{recede}}$) of the approaching and receding lobes are (e.g., Komberg 1994; Ryle & Longair 1967)

$$Q = (1 + \beta \cos \theta)/(1 - \beta \cos \theta), \quad (1)$$

and

$$R = [(1 + \beta \cos \theta)/(1 - \beta \cos \theta)]^{n-\alpha}, \quad (2)$$

with $c\beta$ the bulk velocity of the emitting plasma (assumed for simplicity to be the same on both sides), θ the angle between their motions and the line-of-sight to the observer, α the spectral index (defined so $S_\nu \propto \nu^{+\alpha}$) and $n = 2$ or 3 depending upon whether emission from a continuous flow or discrete blobs, respectively, is assumed. For the hot-spots and lobes, typical values of the spectral index are $-1.0 < \alpha < -0.5$, though some high-redshift sources exhibit steeper spectral indices ($\alpha < -1$).

Since the receding lobe is farther from us, the ages (measured in their rest frames) at which the two lobes are observed are in the ratio

$$t_{\text{recede}}/t_{\text{approach}} = (1 - \beta \cos \theta)/(1 + \beta \cos \theta). \quad (3)$$

After the discovery of compact hot-spots near the outer edges of radio lobes (Hargrave & Ryle 1974; Hargrave & McEllin 1975), it became clear that the asymmetry can be more meaningfully exploited using them (instead of the entire lobe). In that the main thrust of the collimated energy supply from the galactic nucleus is focussed through the jets into the hot-spots, any relativistic motion is more likely to be associated with them (Longair, Ryle & Scheuer 1973; Blandford & Rees

1974; Scheuer 1974), and not with the extended lobes. However, while the flow down the jets may well remain highly relativistic all along, with $\beta_{\text{jet}} \sim 1$ (e.g. Georganopoulos & Kazanas 2004; Urry & Padovani 1995), we shall see that the hot-spot advance speed, β , is much slower.

Numerous studies since then have focussed on the use of the core-hot spot separation ratio as the measure of the ‘asymmetry parameter’, Q , beginning with the early works by Fomalont (1969) and Ingham & Morrison (1975). Assuming an intrinsic source symmetry and a random orientation of the source axis relative to the line-of-sight, these and other authors were able to reproduce the observed distribution of Q among 3CR radio sources (having a median Q of ~ 1.3) with a typical value of $\sim 0.2c - 0.3c$ for the speeds of the hot-spots (e.g., Longair & Riley 1979; Banhatti 1980). Similar speeds were estimated by Macklin (1981), employing the Q -distribution for a well-defined set of 76 3CR sources, and simultaneously fitting the observed distribution of the misalignment angle of the two lobes relative to the nucleus of the parent galaxy. Nonetheless, all these authors concurred that such estimates only provide upper bounds to the true speeds, since part of the apparent asymmetry is likely to be intrinsic to the source. Earlier, Ryle & Longair (1967) had argued that in order for their simple kinematic model, based on the inference of relativistic motion of the lobes, to work, the radio luminosity of the lobes must rapidly ‘evolve’ with time ($P \propto t^{-(3-\alpha)}$). Such a steep temporal fading, however, followed from their assumption that the lobes of double radio sources are a one-time ejection phenomenon.

The first major claim in favor of non-relativistic speeds of the hot-spots came from the work of Mackay (1973); he used the observed distribution of the lobe flux ratios, R , for a well-defined subset of the 3CR complete sample then just mapped with the Cambridge 1-mile synthesis telescope, to set an upper bound of $0.08c$ to the speed of most of the lobes. A prediction of the simple kinematic model is that the approaching lobe should appear not only more distant from the optical identification (due to the light travel time effect), but also somewhat brighter, due to Doppler boosting. However, Mackay (1971) found the opposite: the brighter of the two lobes was seen closer to the optical identification in a vast majority of the 3CR sources. This, so called, ‘Mackay’s rule’ provided a strong hint that double sources recede from the nucleus non-relativistically and that their apparent asymmetry must be of largely intrinsic origin (see, also, Ingham & Morrison 1975). Subsequently, Macklin (1981) was able to demonstrate that usually the hot-spots are more asymmetric in brightness than are the diffuse tails. He further showed that the Mackay’s rule is primarily applicable to the diffuse structure (rather than to the hot-spots), and that this could be a selection effect arising from the heavier expansion losses for the diffuse tail associated with the more distant hot-spot, which presumably resides in a less dense environment (see also, Valtonen 1979). Indeed, a simple way to understand Mackay’s rule would be to imagine that a double radio source is born in an external environment having a density gradient. The lobe moving on the denser side would then be shorter due to larger ram-pressure, but at the same time, brighter, due to a better confinement (Hintzen & Scott 1978; Gopal-Krishna & Wiita 1991; Saikia et al. 1995, 2003; Jeyakumar et al. 2004). Clearly, in this picture the lobe-length asymmetry produced by the dominance of the environmental factor is expected to be weaker for more energetic sources, which again is consistent with observations (e.g., Zięba & Chyży 1991; Saikia et al. 1995, 2003).

Currently, the most popular explanation for the intrinsic structural asymmetry invokes an unequal density of the circumgalactic medium, on the scale of tens of kiloparsecs (e.g., Miley 1980; Swarup & Banhatti 1981; Ensmann & Ulvestad 1984; Fokker 1986; Liu & Pooley 1991b; McCarthy, van Breugel & Kapahi 1991, hereafter, MvBK; Lister, Hutchings & Gower 1994, hereafter, LHG). However, from a new interpretation of the observed correlated radio-optical asymmetry of radio galaxies (MvBK; Liu, Pooley, & Riley 1992), it has been more recently inferred that a substantial part of the (intrinsic) asymmetry may be contributed by the motion of the parent galaxy itself (Gopal-Krishna & Wiita 1996, 2000; §5.1). In either case, the dominance of intrinsic asymmetry

($Q_{\text{int}} \neq 1$) excludes the possibility of applying the elegant formulation of Banhatti (1980) to determine the velocity distribution *analytically* from the observed Q -distribution. It may also be mentioned that several researchers (e.g., Ryś 1994; LHG) have attempted to model the Q -distribution within the framework of the alternate ejection hypothesis proposed by Rudnick & Edgar (1984) (see also, Willis, Wilson & Strom 1978; Wiita & Siah 1981; Komberg 1994), though differing conclusions have been reached (§7.2). The first treatment of the lobe expansion which treated the dynamics of a relativistic hot-spot was given by Gopal-Krishna, Wiita & Saripalli (1989). A recent attempt by Arshakian & Longair (2000) to generalize the Q -distribution analysis of Banhatti (1980) to include both intrinsic and relativistic sources of the asymmetries concludes that they are actually of comparable importance (also see §2.4) for their entire sample of RGs and quasars, but that, perhaps unsurprisingly, the relativistic effects are more important for quasars and the intrinsic ones more so for RGs.

2.2. Subtleties of the asymmetry parameter, Q

As mentioned above, using the 3CR complete sample, Macklin (1981) demonstrated that Mackay’s rule primarily applies to the diffuse tails in the lobes (rather than to the hot-spots), and that this could well be a selection effect. Subsequent studies of the 3CR sample and other samples have led to interesting findings, which are of considerable import for the theories of radio source evolution and their interaction with the environment.

- (i) Radio galaxies become more symmetric with increasing radio luminosity (Zięba & Chyży 1991). However, their asymmetry increases with redshift, z (Zięba & Chyży 1991; MvBK). The former trend would be consistent with the high degree of lobe-length asymmetry ($Q_{\text{median}} \sim 2$) observed in giant radio galaxies (GRGs) which have intermediate radio luminosities and for which projection is least likely to be a significant factor (see Saripalli et al. 1986; Parma et al. 1996; Arshakian & Longair 2000; Ishwara-Chandra & Saikia 2002). It appears that Q , the overall projected linear sizes (ℓ), and the apparent bending angles all evolve with z more strongly for RGs than for quasars (Chyży & Zięba 1995).
- (ii) An early inkling of quasars not following Mackay’s rule came from the work of Riley & Jenkins (1977) on 3CR quasars. More specifically, it was demonstrated by Teerikorpi (1984, 1986) that among quasars, Mackay type asymmetry prevails only up to a certain luminosity, beyond which the sense of asymmetry is reversed. The critical luminosity is $\approx 10^{27} \text{W Hz}^{-1}$ at 1 GHz, taking $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$, and $q_o = 0.5$. His analysis favored the possibility that high-luminosity quasars (rather than high- z quasars) tend to disobey Mackay’s rule. With hindsight, this is what would be expected, since it is in more luminous quasars that the light-travel-time effects are expected to be large, due to a combination of their hot-spot moving with a higher speed and in a direction relatively close to the line-of-sight (within the framework of the ‘unified scheme’; §2.3). This point is further discussed below.
- (iii) For quasars, there seems to be no strong dependence of Q on redshift, at least up to $z \sim 2$ (e.g., Kapahi 1990a; also, LHG). In contrast, for 3CR RGs, MvBK have noticed a statistical tendency for Q to increase with redshift, over the range $0 < z < 2.5$, as also suggested by Zięba & Chyży (1991). Regardless of whether the lobe-length asymmetry arises from environmental asymmetry/inhomogeneity (§5), or quite substantially from the motion of the parent galaxy itself (§5.1; Gopal-Krishna & Wiita 1996), the increased asymmetry found for galaxies at high redshifts would be expected to be manifested through a corresponding rise in the apparent bending angle, Ψ , (a measure of departure from colinearity of the twin lobes with the nucleus), particularly since at higher redshifts the physical sizes of radio sources were smaller (Kapahi 1989; Singal 1988). Indeed, a comparison of radio structures of lobe-dominated quasars at $z \leq 1.5$ and $z \geq 1.5$ led Barthel

& Miley (1988) to suggest that quasars are more bent at higher redshifts. However, this oft-cited claim was questioned by Kapahi (1990b) who failed to find a strong effect, in spite of using maps that were better matched in *linear* resolution. Settling this issue is of considerable significance for cosmology. Probably a more appropriate approach would be to employ samples of RGs alone (as also used for investigating the $Q - z$ dependence), because their structures are believed to be less distorted by projection effects, and they are now being found in substantial numbers at high redshifts.

2.3 Lobe-length asymmetry in the context of the Unified Scheme for powerful radio sources

According to a widely discussed unified scheme for powerful radio galaxies and quasars, the latter are RGs with the AGN axis oriented within a certain critical angle ($\theta_c \approx 45^\circ$) from the line-of-sight (Barthel 1989; also, Scheuer 1987; Peacock 1987; Morisawa & Takahara 1987; Antonucci & Barvainis 1990). In an early examination of orientation effects, a correlation between the two orientation dependent parameters, Q and the core-fraction, f_c (the flux ratio of the radio core flux to total flux) was searched for using a set of 26 3CR sources with detected radio cores (Gopal-Krishna 1980). Subsequently, using larger samples of quasars, a weak but significant Q - f_c correlation was found (e.g., Kapahi & Saikia 1982; Saikia 1984; Hough & Readhead 1989), though the effect has not been discerned in some other quasar samples (e.g., Lister, Gower & Hutchings 1994). Baryshev & Teerikorpi (1995) have analyzed the asymmetry properties of double RGs within the context of the unified scheme. They allowed for intrinsic asymmetries, including non-zero misalignment angles, and performed numerical studies to bolster their analytical models for velocities and other properties of quasars and RGs. More recently, Saikia et al. (1995, 2003) have found evidence for a higher degree of asymmetry among compact-steep-spectrum (CSS) quasars, *vis-à-vis* CSS RGs, again in broad agreement with the orientation-based unified scheme. Such asymmetries are reproducible through propagation of symmetric jets through asymmetric confining media as shown by the analytical models and numerical simulations of Jeyakumar et al. (2004).

One important check on the consistency of the Q -distributions with the unified scheme comes from the work of Best et al. (1995), which employs the well observed 3CRR complete sample. For a subset of 95 FR II sources with a well-identified hot spot in each lobe, they found that the quasars appear distinctly more asymmetric than the radio galaxies and that this difference is not due to any luminosity or redshift bias. Likewise, they also found that, compared to galaxies, the radio lobe-pair in quasars appear more misaligned (i.e., less colinear with the nucleus). Note that, unlike Q , the apparent misalignment angle (Ψ) of the twin-lobes does not depend upon the hot-spot speed. Taking the critical viewing angle (θ_c), which separates galaxies from quasars in the unified scheme, to be 45° (Barthel 1989), Best et al. (1995) deduced that typically the *intrinsic* misalignment angle of the hot-spots is randomly distributed within a cone angle $\phi_{max} = 10^\circ$. Adopting these values of ϕ_{max} and θ_c , they could then reproduce the observed marked difference between the Q -distributions for the RGs and quasars, taking typical hot-spot speeds of $0.2c - 0.3c$. Further, making an allowance for an (environmentally induced) intrinsic asymmetry, they deduced a marginally lower value of $\sim 0.16c$. Similar estimates had earlier been reported by LHG, from their orientation-based modeling of the radio structural parameters for a large (but not complete) sample of 192 powerful RGs and quasars.

While such large hot-spot speeds conform to the estimates coming from the analysis of synchrotron ageing effects in the radio lobes in powerful sources at $z \geq 0.2$ (e.g., Liu, Pooley & Riley 1992; also, Leahy, Muxlow & Stephens 1989), an analysis of the Q -distributions of quasars by Scheuer (1995), within the framework of the unified theory (which imposes a θ_c), has indicated that even for the most powerful sources, the hot-spot speeds are only a few percent of the speed of

light. As discussed below (§2.4), the extra clue employed by Scheuer is the apparent jet-sidedness, which can be used to identify the approaching and receding lobes (see, also, Saikia 1984). Thus, the origin of the substantially higher values of β obtained in the analysis of Best et al. (1995) remain to be fully understood, particularly in view of their assertion that such high speeds are necessary to explain the observed contrast between the distributions of Q and Ψ for powerful RGs and quasars, within the framework of the unified scheme.

The differences in cosmological evolution of ℓ between RGs and quasars, as well as the stronger correlations of ℓ with radio luminosity for RGs than for quasars, may imply difficulties for the pure orientation based unified scheme (Chyży & Zięba 1995). We note, however, that these differences can be readily understood within a more realistic unified scheme which takes into account the growth of radio sources with time as well as a rise in θ_c with intrinsic source power (Gopal-Krishna, Kulkarni & Wiita 1996; also, Lawrence 1991).

2.4 Further constraining the hot-spot speeds by the jet-sidedness

As mentioned above, Scheuer (1995) has argued that a much sharper interpretation of the Q -distribution is possible by invoking additional information, namely, the presence of the jet, which allows one to distinguish the approaching lobe/hot-spot from its receding counterpart. This clue was earlier employed by Saikia (1984), but his sample of 36 quasars did not show the expected trend; namely a statistical excess of the lobe-length on the jet side. Since the orientation dominated effects are likely to be stronger for highly luminous radio sources with their hot-spots advancing more rapidly, Scheuer has determined the lobe-length ratios for 3 essentially independent samples containing a total of 43 sources with conspicuous jets (including 41 quasars), above an emitted integrated luminosity of $\sim 10^{27.5} \text{ W Hz}^{-1} \text{ sr}^{-1}$ at 178 MHz. For each of these high luminosity samples, he finds the jet-side lobe to be systematically, albeit only marginally, longer than the opposite (receding) lobe. This is basically in agreement with the kinematical model. Assuming that θ is uniformly distributed over solid angle between $\theta = 0$ and $\theta = \theta_{\max}$, the predicted median of Q and the mean of the logarithm of Q are (Scheuer 1995):

$$Q_{\text{median}} = \frac{1 + 0.5\beta(1 + \cos\theta_{\max})}{1 - 0.5\beta(1 - \cos\theta_{\max})} = \beta(1 + \cos\theta_{\max})[1 + \frac{1}{3}\beta^2(1 + \cos\theta_{\max})^2 \dots], \quad (4)$$

$$\langle \ln Q \rangle = \int_{\theta=0}^{\theta_{\max}} \ln Q \, dP = \frac{f(\beta) - f(\beta \cos\theta_{\max})}{\beta(1 - \cos\theta_{\max})}, \quad (5)$$

where $f(\beta) = (1 + \beta) \ln(1 + \beta) + (1 - \beta) \ln(1 - \beta)$.

Comparison of the observed Q -distributions with the predicted ones yields the fairly robust results: (i) $\beta \geq 0$, and (ii) $\beta \leq 0.15$, with the most probable value being $\beta = 0.03 \pm 0.02$ (Scheuer 1995). It is remarkable that these estimates of β from the source asymmetry (which, obviously, are independent of the adopted values of the cosmological parameters H_0 , Ω and Λ) are well below the corresponding value of $\beta \sim 0.165$ deduced by Scheuer from the synchrotron ageing results reported by Liu, Pooley & Riley (1992). The substantial difference between the two estimates was taken by Scheuer as evidence for a rapid ‘backflow’ of the relativistic plasma within the lobes, which is indeed expected if the jet material is much less dense than the ambient plasma. Such strong backflows are clearly observed in many numerical simulations of jet propagation (e.g., Norman 1996; Hooda & Wiita 1998).

It is worth stressing that even though the synchrotron ageing estimates are still routinely used, they have been criticized from several other different angles (e.g. Wiita & Gopal-Krishna 1990; Siah

& Wiita 1990; Rudnick, Katz-Stone & Anderson 1994; Eilek & Arendt 1996; Blundell & Rawlings 2000). See §5.2 for a more detailed discussion of radio galaxy ages.

The most recent analysis of high quality data leading to a determination of β for the hot-spots is due to Arshakian & Longair (2000). They also use the jet-sidedness ratio, which they could estimate for 80 per cent of the FR II sources in the 3CRR complete sample, giving them a total of 103 sources (71 RGs and 32 quasars). They conclude that the asymmetry is dominated by relativistic effects for the quasars, in the sense that the approaching jet is typically on the long lobe side. But just the opposite is the case for RGs, where there are more sources with approaching jets on the short lobe side; this is indicative of the intrinsic asymmetry (or host motion asymmetry) for these objects. Their best estimate of the mean hot-spot velocities is $\langle\beta\rangle = 0.11 \pm 0.013$, substantially above Scheuer's, and not too dissimilar from the results estimated using the unreliable spectral ageing technique. Within the framework of a Gaussian model, the standard deviation of the velocity distribution is found to be $\sigma_\beta = 0.04$. Arshakian & Longair (2000) note that these results are in agreement with the unified model with $\theta_c \simeq 45^\circ$. Although this work is rather convincing, it is clear that substantially larger samples of RGs and quasars with well defined jet-sidedness will be needed to finally settle the question of the average value of β and if it evolves with P , z , and/or ℓ .

2.5. Brightness asymmetry of hot-spots

Besides the asymmetric locations of the hot-spots relative to the core (§§2.1–2.2), another striking asymmetry related to the hot-spots has recently been noticed. This new result concerns the brightness asymmetry of the hot-spots, which is specially pronounced in the X-ray emission from the hot-spots of quasars and broad line radio galaxies.

Although just a handful of such examples are known at present, the trend seems quite clear that the brighter of the two hot-spots in such sources lies on the side of the main jet (Georganopoulos & Kazanas 2003). These authors argue that this X-ray asymmetry arises from the inverse Compton boosting of the radio photons produced inside the hotspot, where the jet plasma undergoes a rapid deceleration, resulting in the emission of inverse Compton X-rays which is beamed more narrowly than is the nonthermal radiation at lower frequencies. The hotspot asymmetry caused by such an energy dependent beaming furnishes an important piece of evidence for the idea that bulk motion of the jet remains relativistic all the way to the hot-spots (Georganopoulos & Kazanas 2003). With the increasing detections of hot-spots with *Chandra*, rapid progress in this area can be expected.

3. The asymmetric depolarization of the twin radio lobes

The anomalously large difference between the Faraday rotation measure of the radio lobe pair was first discussed for the case of Cygnus A, and was interpreted in terms of differential Faraday depth to the two lobes (Slysh 1966). In 1988, a striking correlation, now called the ‘Laing-Garrington (L-G) effect’ was demonstrated using a sample of double-lobed radio sources (mostly quasars) showing one-sided jets. According to the correlation, the lobe on the jet-side depolarizes less rapidly with increasing wavelength than does the opposite lobe (Laing 1988; Garrington et al. 1988).

The currently popular explanation of this L-G effect, advanced by its discoverers, posits that distant radio sources are embedded within ≈ 100 kpc diameter halos of hot gas, similar to the cores of the intra-cluster medium (ICM) associated with nearby clusters of galaxies. Consequently, the lobe on the near-side, which is also observed to be associated with the (presumably Doppler-boosted) extended jet, is viewed through a smaller magneto-ionic depth of the halo and hence suffers less depolarization (see, also, Tribble 1992). For quasars, whose axes are believed to be oriented away from the plane of the sky, it is reassuring that the depolarization asymmetry is indeed found to be

governed by the jet-sidedness, rather than by any difference in the lobe-lengths (Garrington, Conway & Leahy 1991). Furthermore, for RGs, for which the differential magneto-ionic depth to the two lobes would normally be a less crucial factor (e.g., Barthel 1989), it is the shorter lobes which tend to suffer greater depolarization. This can be readily understood as a result of radially declining density profile of the halo plasma (Garrington & Conway 1991; Tribble 1992; Pedelty et al. 1989; also, Strom & Jägers 1988).

Although the above explanation for the L-G effect in (distant) quasars, invoking an ICM-core like ambient medium, has almost become a folklore, a few disconcerting aspects have been highlighted by Gopal-Krishna & Nath (1997), leading them to consider an alternative scenario. Firstly, the viability of the ‘standard’ explanation rests on the tacit assumption that in every case, the postulated ICM core around the radio source is somehow able to maintain a diameter very close to the steadily growing size of the source. Even a factor of two discrepancy between the two sizes would seriously erode the correlation (Garrington & Conway 1991). However, no regulatory mechanism has been suggested that would ensure the needed tight coupling between the two sizes. Secondly, whilst the L-G effect is found to become stronger at higher redshifts, the density, magnetic field, structure, and even the existence, of a hot ICM core around all high- z quasars is quite uncertain. In the absence of such deep, high resolution X-ray observations, the basic assumption of the canonical model, namely, the general existence of a dense ICM core ($n > 10^{-2}\text{cm}^{-3}$) around $z > 1$ quasars (Garrington & Conway 1991) is fraught with uncertainty (though, clearly, even distant radio sources must still have a low-density gas envelope).

3.1. An alternative scenario for the origin of depolarization asymmetry

In view of the above caveats, it has been proposed that the required Faraday screen contributing the extra rotation measure (RM) toward the far-side lobe is associated not with any putative ICM, but, instead, with a large disk of magneto-ionic medium surrounding the host galaxy and oriented roughly perpendicular to the radio source axis (Gopal-Krishna & Nath 1997; Gopal-Krishna & Wiita 2000). As discussed below, there is a growing evidence for large disks of dust and gas around RGs and quasars, even at high redshifts where the L-G effect is found to be strong. Such dusty *superdisks*, or *fat pancakes* could be responsible for a variety of phenomena associated with radio sources (§5). Their prominent tracers, on smaller scale, are the extended dust-lanes which tend to align perpendicular to the jets (Kotanyi & Ekers 1979). They have been detected in roughly half of the bright ellipticals, despite the difficulty of detection in the absence of a fortuitously edge-on view (Goudfrooij & de Jong 1995 and references therein). At large redshifts ($z > 2$), evidence for such dusty *superdisks* around RGs and quasars comes from the apparent suppression of the Ly- α emission associated with the far-side lobe of radio galaxies (Gopal-Krishna et al. 1995) and quasars (Heckman et al. 1991a) (§6). According to a widely discussed scenario, dusty disks around the elliptical hosts of powerful radio sources form as a result of accretion of gas-rich galaxies, which also provides a trigger for the nuclear activity (e.g., Fukugita, Hogan & Peebles 1996; Lynden-Bell 1996; Wilson & Colbert 1995; Colina & de Juan 1995; also, Sofue & Wakamatsu 1992). In fact, optical observations have revealed an abundant population of gas-rich field galaxies undergoing massive starbursts at redshifts down to $z \sim 1$ (e.g. Cowie, Hu & Songaila 1995). Alternatively, these superdisks could be parts of the cosmic web of filamentary network of proto-galactic material, as revealed in a number of simulations of the structure formation process (e.g., Cen & Ostriker 1999).

Independent evidence for the putative *superdisks* around radio galaxies has emerged from observations in the radio band (see Gopal-Krishna & Wiita 2000 for a more comprehensive review). For instance, millimetric continuum emission believed to arise from an enormously massive dusty disk ($M_{\text{gas}} \sim 10^{11-12} M_{\odot}$) has been detected in the extremely distant ($z = 3.8$) radio galaxy, 4C41.17

(Dunlop et al. 1994; Chini & Krügel 1994). Striking evidence for fat and highly extended gaseous disks around the elliptical hosts of RGs comes from the detection of sharp and often quasi-linear inner boundaries of the radio lobes, creating strip-like central gaps in the radio bridges (Gopal-Krishna & Wiita 1996; 2000). Although their detection requires not only maps with high sensitivity and dynamic range, but also a favorable orientation of the lobe axis (i.e., close to the plane of the sky), clear examples of such sharp inner boundaries of radio lobes have been observed in a number of RGs. These include: 3C34 (Johnson, Leahy & Garrington 1995); 3C227 (Black et al. 1992), 0828+32 and 3C382 (Capetti et al. 1993) and 3C16, 3C33, 3C61.1, 3C184.1, 3C341, 3C381, 4C14.11 and 4C14.27 (Leahy & Perley 1991). An example of such morphology is shown in Fig. 1.

The sharp-edged central emission gaps can be attributed to the blocking of ‘back-flowing’ non-thermal plasma of the radio lobes by a denser thermal plasma associated with the parent galaxy. Thus, from the ‘height’ of the radio gaps in these 12 sources ($z_{\text{median}} \sim 0.2$) we estimated a median value of ≈ 75 kpc for the disk diameter, which is clearly a lower limit. Likewise, the observed ‘widths’ of the gaps suggest a median value for the disk thickness, $l_{\text{disk}} \approx 25$ kpc. Such enormous disks (*superdisks*) are consistent with the numerical simulations of the galaxy formation at high redshifts in hierarchical clustering scenarios (e.g., Evrard, Summers & Davis 1994). Since quasars at $z > 1$ have a median radio size of $\sim 60\text{--}70$ kpc (taking $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $q_0 = 0.5$), *their far-side lobes could well be largely hidden behind the postulated superdisks.*

Gopal-Krishna & Nath (1997) have discussed a scenario for the thermal evolution of the tidally-stretched gaseous disk of the captured galaxy, as it gravitates further toward the interior of the hot gaseous corona of the massive elliptical. They estimate that for plausible gas densities, the disk would continue to tap heat from the ambient coronal gas and expand in width to ≈ 10 kpc in $10^8\text{--}10^9$ yr, which is the expected formation timescale for the dust-lanes. Such widths are consistent with the observed dimensions of the gaps along the radio bridges. Interestingly, direct evidence for depletion of cool interstellar medium (HI) of a captured disk galaxy, *via* heat input from the coronal gas of the captor galaxy, has come from HI imaging of the early-type galaxy NGC1291 (Bregman, Hogg & Roberts 1995). Note also that in addition to widening it, the heat input into the disk is also expected to steadily increase the diameter of its ionized central part, concurrently with the growing size of the radio source. The estimated average disk parameters (magnetic field, $B \sim 1 \mu\text{G}$, and electron column density, $\Sigma \simeq 300 \text{ pc cm}^{-3}$) correspond to a Faraday dispersion parameter, $\Delta \approx \langle n^2 B^2 \rangle^{1/2} (l_{\text{disk}})^{1/2} \simeq 10^2 \text{ cm}^{-3} \mu\text{G pc}$ (Gopal-Krishna & Nath 1997; see also Gopal-Krishna & Wiita 2000 – hereafter GW00). Such values are quite consistent with those inferred from the depolarization maps of quasars (Garrington & Conway 1991). Sensitive UV/X-ray imaging with space observatories could possibly provide strong support of this scenario, via a direct detection of the hot component of the postulated magneto-ionic superdisks around powerful, distant radio galaxies, although the combination of spatial resolution and sensitivity seems to be just beyond the current generation of X-ray telescopes (*Chandra* and *XMM-Newton*). Perhaps a more promising strategy to detect such disks would be to map the dust-reddening of the background galaxies.

4. The radio spectral-index asymmetry

While the depolarization asymmetry (Laing-Garrington effect) can be naturally explained in terms of orientation effects (§3), there is still no consensus on the origin of another prominent asymmetry which involves the radio spectral index. This intriguing (statistical) asymmetry was noticed in two separate studies of 3CR radio sources, and can be summed up as follows:

- (a) The lobe which is less depolarized has a flatter radio spectrum (Liu & Pooley 1991b).
- (b) The lobe on the jet-side has a systematically flatter spectral index by typically $\Delta\alpha \sim 0.1$, though with a large dispersion (Garrington et al. 1991).



Figure 1: Total intensity radiograph for 3C33 at 2 cm, made with the VLA, showing the sharp inner boundaries of the lobes. Reprinted with permission from Leahy & Perley, *AJ*, 102, 537 (1991). ©American Astronomical Society

Note that the first statement refers to a sample consisting chiefly of radio galaxies, for which orientation effects are expected to be weak, though not necessarily negligible. Specifically, for such sources the longer lobe tends to be less depolarized, which is consistent with the halo of magneto-ionic plasma having a decreasing radial profile (e.g., Pedelty et al. 1989; also, Strom & Jägers 1988).

As argued by Liu & Pooley (1991a,b), the effect (a) does not arise from the jet contaminating the approaching lobe’s emission, since the jet contribution would be inadequate. These authors also discount the possibility that pure Doppler shifts, in conjunction with a downward curved spectrum of the radio lobes could explain the observed spectral asymmetry (see, also, Blundell & Alexander 1994). Nonetheless, Doppler effects could lead to strong flux boosting of the approaching hot-spot, thus raising its contribution to the flux density of the nearer lobe (which is associated with the observed jet). This is also consistent with the result that the hot-spot on the jet side is usually brighter (Bridle & Perley 1984; Laing 1988; Teerikorpi 1984; Dennett-Thorpe et al. 1997). Since the hot-spot spectrum is flatter than the lobe spectrum, the enhanced hot-spot contribution would flatten the integrated spectrum of the nearer lobe (which is associated with the brighter jet and is usually less depolarized), as compared to the farther lobe (Garrington et al. 1991). Extending this argument, Tribble (1992) estimated that the approaching hot-spot *itself* might exhibit a measurably flatter spectrum compared to the receding hot-spot, since the core of the approaching hot-spot can be expected to have not only a flatter spectrum than the annular portion, but also a greater Doppler boost due to a faster forward motion (which, incidentally, could be well above the time-averaged speed of the hot-spot; Scheuer 1982). One may thus be able to understand not just the compactness asymmetry of the hot-spots (Laing 1995; see Tribble 1992) but also their spectral-index asymmetry, as reported by Liu & Pooley (1991a,b). Tribble has further argued that the rather strong correlation between the depolarization asymmetry and the spectral-index asymmetry is an indication that the latter, too, is largely an orientation effect.

4.1. A further twist

However, another study calls for caution in accepting the apparently attractive explanation of the spectral-index asymmetry of the lobes in terms of a greater contamination from the brighter, Doppler-boosted hot-spot present on the jet side (see above). From detailed VLA observations of a set of 8 powerful radio quasars, Bridle et al. (1994) have shown that nearly in every case, the hot-spot on the jet side has *itself* a flatter spectrum than the hot-spot on the counter-jet side. Moreover, the hot-spots on the jets’ sides do not appear systematically brighter. While it is important to settle these points conclusively by mapping a fairly large sample of quasars with the VLBA, a few interesting possibilities emerge from the work of Bridle et al. Basically, they find that whereas the radio spectrum of high surface brightness features (hot-spots) is almost always flatter on the jet-side, the diffuse lobe emission on that side has a *steeper* spectrum, if the associated lobe is significantly shorter than the other lobe. If true, this could imply a link between the mechanisms of producing a shorter lobe on the jet side (which is unexpected in the simple kinematical model; §2.1) and the energy loss rate of the radiating electrons within the lobes. It is clearly important to establish the generality of the result of Bridle et al. (1994), which suggests that while the jet-side controls the spectral asymmetry of the bright compact emission, the spectral asymmetry of the diffuse emission is controlled by the lobe-length asymmetry.

An ‘intrinsic’ explanation for the spectral asymmetry of the hot-spots/lobes on the two sides has been proposed in terms of a steady dilution of the magnetic field as the radio source grows, essentially in a self-similar fashion (Blundell & Alexander 1994). From a quantitative, albeit highly model-dependent, treatment based on this idea, these authors are able to explain the typically observed spectral asymmetry ($0 < \Delta\alpha < 0.4$). They argue that, due to projection and differential light

travel time effects, the lobe on the counter-jet side is monitored at an earlier stage, when its magnetic field is postulated to have been stronger and hence its spectrum steeper, thanks to heavier radiative losses. This simple scenario predicts: (i) a higher surface brightness for the lobe with a steeper spectrum and stronger depolarization; and (ii) the magnitude of the spectral asymmetry should be maximum for the outer edges of the lobes and minimum for the parts close to the nucleus. Subsequent observations by Dennett-Thorpe et al. (1997) did not bear out this prediction, as they found that for more aligned jets, the near hot-spot has a higher flux and a flatter spectrum; however, their sample is small (just 6 sources with detectable jets). Interestingly, these authors find that in 5 of the 6 sources the lobe with a larger surface area has a flatter spectrum. Thus, at this stage, it remains a challenge to disentangle the intrinsic, environmental and orientational effects in the interpretation of the spectral asymmetry, and only larger samples will allow these important tasks to be accomplished.

4.2. Correlation of the structural asymmetry with spectral curvature

Yet another type of asymmetry was noticed from a careful analysis of the radio spectra of a well-defined large sample of FR II radio galaxies selected from 3CR sources. The available data for these galaxies allowed the determination of the *rest-frame radio spectrum* for individual sources, after subtracting the compact radio emission associated with the core. It was thus found that, statistically, the lobe spectra are more curved for sources with larger core-to-hot spot separation ratio (Q) (Mangalam & Gopal-Krishna 1995). This correlation between spectral-curvature (C) and asymmetry (Q) needs to be further investigated using a larger sample, as it can provide potentially useful clues about the evolutionary tracks for powerful radio sources.

It is instructive to consider the Q - C correlation in conjunction with the result that a larger spectral curvature is more prevalent among FR II sources having normal spectra (i.e., not very steep) near the rest-frame frequency of 1 GHz (Mangalam & Gopal-Krishna 1995). This is suggestive of a temporal evolution heading toward not only a steepened but also a straightened spectrum (within the observed radio window), saturating near $\alpha \approx -1.2$ (Mangalam & Gopal-Krishna 1995). Since such an evolution of powerful radio sources could arise from an accumulation of energy losses, the source asymmetry should be linked to this factor, in view of the Q - C correlation. One possibility is that within a radius of ≈ 10 kpc of the nucleus the circum-galactic medium is not only dense but also very clumpy. At least for distant RGs at $z \geq 2$, an abundance of cool gaseous clumps in the circum-galactic medium has been inferred from the observed multiple absorption troughs in the extended Lyman- α emission profiles of RGs with dimensions up to ~ 50 kpc (van Ojik et al. 1996). The presence of such a clumpy circum-galactic medium can have a profound influence on the early evolution of radio sources, leading to a conspicuous structural asymmetry and, concurrently, a curved radio spectrum, by slowing down the jets and thereby prolonging the efficient confinement of the lobes (Mangalam & Gopal-Krishna 1995; Saikia et al. 1995; Wang, Wiita & Hooda 2000; Mellema, Kurk & Röttgering 2002).

An interesting possible link between the structural asymmetry and the presence of a dense ISM in the parent galaxy is hinted by the work of Cimatti et al. (1993). They found a tendency of markedly asymmetric radio sources to show substantially polarized light across the parent galaxy; the polarization probably results from scattering of the nuclear light by a dense ISM of the galaxy.

5. Correlated radio-optical asymmetries in powerful radio galaxies

A major impetus for linking the environmental factor to the asymmetries has come from the study of the extended optical emission-line regions (EELR) associated with the FR II type 3CR radio galaxies, as reported by McCarthy & van Breugel (1989); MvBK (see, also, Liu & Pooley 1991b).

These authors found a *very strong* tendency for the optical surface brightness to peak on the side of the shorter radio lobe. Further, the optical brightness contrast between the two sides was found to be rather pronounced for higher redshift sources ($z > 0.3$) for which the line emission was measured in terms of the blue line [O II] λ 3727. MvBK proposed that the causal link between these tightly correlated radio and optical asymmetries is a large-scale density asymmetry in the circum-galactic medium of the radio-loud elliptical, which slows the lobe propagation more effectively on the denser side (e.g., Hintzen & Scott 1978; Swarup & Banhatti 1982), while simultaneously enhancing the optical output of the thermal filaments of the EELR on that same side *via* a stronger confinement of the filaments.

Although appealing for its simplicity, this hypothesis invoking environmental asymmetry does not seem to offer a robust explanation for the *nearly universal* correlation which is found to hold even when the lobe-length asymmetry is at a level of just $\approx 1\%$ (MvBK), particularly in view of the fact that in a given lobe the physical separations of the radio and optical peaks from the nucleus usually differ by an order-of-magnitude (MvBK). On the other hand, should the spatial scale of the environmental density asymmetry indeed be large enough to encompass the entire radio lobe, the putative excess ambient pressure enhancing the optical line-emission from the shorter lobe through a more effective confinement on that side should also enhance that lobe’s radio output. No significant trend of this sort is noticed, however (MvBK). Furthermore, it remains unclear why the optical asymmetry is more pronounced for higher redshift galaxies whose EELRs were imaged in the blue [O II] λ 3727 line, as compared to the low-redshift galaxies imaged in the (red) H α line (MvBK). Note also that if the trajectories of the twin jets are not precisely anti-parallel, or linear (which is likely, given the “dentist’s drill” behavior of the hot-spots; e.g., Scheuer 1982; Clarke 1996; Norman 1996), then the asymmetry parameter, Q , could easily depart mildly from unity, without there being any systematic difference present between the two lobes’ environments. These various difficulties have led us to conclude that invoking a nearly universal environmental density asymmetry about radio-loud ellipticals, in order to explain the tight radio-optical correlated asymmetry is probably too simplistic, and a *more foolproof* mechanism is likely to be at work (Gopal-Krishna & Wiita 1996; hereafter GW96).

5.1. A new scenario for the correlated radio-optical asymmetry and the lobe-length asymmetry

Our proposal is based on the premise that the motion of the parent galaxy against the external medium will cause a ram pressure sweeping of the (dusty) ISM, leading to its asymmetric disposition about the galaxy. Consequently, the dust obscuration of the nuclear UV photons and of the light emitted from the EELR filaments will become stronger and more widespread on the downstream side of the galactic center. At the same time, the radio lobe on that side would be automatically identified as the longer one, due to the continuing motion of the galaxy in the opposite direction. Not only would this naturally engender the correlated radio-optical asymmetry, but it would also explain the strong tendency for the EELR asymmetry to be more marked at higher redshifts, since the observed optical window would then sample shorter rest-frame wavelengths, where the dust obscuration is expected to be more pronounced. Details of this scheme are spelled out in GW96 (see, also, §5.3).

Using a sample of 12 radio galaxies showing evidence for a superdisk, GW00 noted an interesting trend: the lobe-length asymmetry is systematically milder when the lobe lengths are measured from the mid-plane of the radio emission gap (the superdisk), instead of employing the usual definition when the lobe lengths are measured using the radio nucleus as the reference point. The offset of the radio nucleus from the superdisk midplane thus indicates a motion of the host galaxy relative to the

superdisk, which is possibly a portion of a cosmic filament. Thus, within a typical age of a radio source, $\tau > 10^8 \text{ yr}$ (§5.2), the parent galaxy would have moved a distance $D = 30\tau_8 V_{g(300)} \text{ kpc}$ from the origin, marked by the point of symmetry between the two hot-spots, where the lifetime τ_8 is in units of 10^8 years and the galactic velocity component along the radio axis, $V_{g(300)}$, is in units of 300 km s^{-1} . Based on the results of various published simulations, GW96 adopted $V_{g(300)} \sim 1$ for the powerful radio sources at $z > 0.5$, which tend to occur inside clusters of galaxies (Yee & Ellingson 1993, and references therein). Now, a D of $\sim 30 \text{ kpc}$ corresponds to an asymmetry parameter $Q \sim 2$, given that the typical size of powerful RGs at $z > 0.5$ is $\sim 150 \text{ kpc}$. Since, for distant RGs the observed $Q_{\text{median}} \sim 1.5$, it seems that the motion of the parent galaxy should play a very significant role in causing the lobe-length asymmetry, though some rare, highly asymmetric, sources would still warrant alternative explanation(s).

5.2. Estimated lifetime of nuclear activity in double radio sources

Historically, the ages of the radio sources have been determined from the observed large-scale spectral gradients along the lobe’s length, by taking into account the main energy loss processes, involving radiative and expansion losses of the energetic particles. The ages thus derived are typically of order of 10^7 years (e.g., Leahy, Muxlow & Stephens 1989; Liu et al. 1992). However, a few authors have questioned the various assumptions underlying this method (e.g., Rudnick et al. 1994; Blundell & Rawlings 2000, 2001; see Wiita & Gopal-Krishna 1990, for one of the earliest criticisms of the spectral ageing method). For instance, in their critique, Blundell & Rawlings (2001) have argued that the ages may have been grossly underestimated on account of the following (unrealistic) assumptions: 1) temporal constancy of magnetic field in the lobes; 2) power-law nature of the electron injection spectrum into the lobes; 3) radiative lifetimes of the synchrotron emitting particles in the lobes are longer than the ages to be deduced; 4) no mixing of the relativistic particle population within the lobes.

By modelling a several flux-limited samples, totalling 303 radio sources selected near 0.15 GHz, their best-fit to the data in the spectral index–size–luminosity–redshift ($\alpha - P - D - z$) space yielded a typical activity lifetime of 5×10^8 years (Blundell, Rawlings & Willott 1999). These longer lifetimes are in better accord with the ages inferred from the hot spot velocities estimated from lobe length ratios (e.g., Scheuer 1995; §2.4). Because of severe inverse Compton losses to the microwave background at high redshifts, very few large, therefore old, sources are detectable at higher z (Gopal-Krishna et al. 1989); this was an early version of the so-called “youth-redshift degeneracy” (Blundell et al. 1999).

More recently, independent support for such long nuclear activity has come from several studies made in other wavebands, and we now note a few of these investigations. Employing very deep VLA observations of a hard X-ray selected sample of 69 sources, Barger et al. (2001) have found that almost 4% of the $L > L_*$ galaxies are X-ray luminous at any time, indicating an average activity lifetime of $\sim 5 \times 10^8$ years for their supermassive black holes. Using a large sample of ~ 5000 massive galaxies at $z < 0.1$, detected in the SDSS Survey (York et al. 2000), Miller et al. (2003) have found that almost 40% of them contain an AGN. From the high fraction of AGN among the massive black holes in the local universe, they infer an average activity lifetime to be $> 10^8 \text{ yr}$. Most recently, Marconi et al. (2004) have estimated the local ($z = 0$) distribution of black hole mass to be $\sim 5 \times 10^5 M_\odot \text{ Mpc}^{-3}$. Interpreting these as exclusively the relics of AGN activity (since $z < 3$), they estimate an average total activity lifetime to be in the range $10^8 - 10^9$ years.

These new data supporting long periods of RG activity add additional weight to our argument that galaxy motion is likely to produce a substantial portion of the observed asymmetries (§5.1).

5.3. Asymmetric obscuration, instead of asymmetric emission

While careful optical/UV observations have strengthened the case for distributed dust within the ISM of RGs and quasars (e.g., Goudfrooij & de Jong 1995; Wills et al. 1993), evidence is also emerging for extended dusty superdisks in such objects, as summarized in §3.1. The latter can act as a rich reservoir of dust which is gradually released into the diffuse ISM on the downstream side, as the galaxy moves on. GW96 have discussed evidence for the galaxy motion, based on the observed offsets of the radio nucleus from the mid-point of the radio emission gap perceived to coincide with the dusty disk. As argued in GW96 and GW00, the combined effect of the dust sweeping from the ISM and the disk component into the downstream side of the moving host galaxy would be to enhance the degree and domain of obscuration on that side *vis-à-vis* the upstream side which would *necessarily* coincide with the shorter radio lobe. The result would be a diminishing of the magneto-ionic content, and especially a decline in the EELR emission, on the side of the longer lobe. This is expected, in that an enhanced dust attenuation of the nuclear ionizing photon flux would be compounded by the attenuation of the UV/blue light emerging from the EELR.

To summarize, in our picture, the tightly correlated radio-optical asymmetry of powerful RGs is not contingent upon the conventional assumption of a nearly universal density asymmetry about the host galaxy. Instead, the correlated asymmetry is driven largely by the galaxy motion which leads to asymmetric obscuration on the two sides. The various difficulties with the conventional explanation can thus be circumvented (see above). These include: (i) the radio-optical asymmetry correlation being present even if the lobe-length ratio departs only minutely from unity, in spite of the radio and optical emission originating on vastly different physical scales; (ii) the asymmetry being much more pronounced in the (blue) line [O II] λ 3727, as compared to the (red) H α line; and (iii) the tendency of the shorter lobe to be frequently the fainter one in the radio, though the opposite would always be expected in case the lobe-length asymmetry were firmly rooted in an environmental asymmetry. Variants of this idea and additional implications are discussed in GW00.

6. Lyman- α halos as the diagnostic of dusty superdisks in high- z RGs

A prime motivation to search for galaxies at high redshifts has been the prospect of detecting their extended emission components which, in the case of (distant) quasars, are usually swamped by the bright (point-like) nuclear emission. Furthermore, the giant Ly- α nebulosities found associated with many distant RGs can provide a sensitive probe of dust in the early universe, since Ly- α photons, being resonantly trapped within the nebula, can be efficiently absorbed by even a small amount of dust. Moreover, as discussed by Gopal-Krishna et al. (1995) and GW00, any conspicuous brightness asymmetry of the Ly- α nebulosity about the galactic nucleus could often be useful for distinguishing the near-side lobe from its counterpart on the far-side of the nucleus; making this distinction normally remains only a cherished dream, since radio jets often remain undetected in powerful RGs.

Fig. 2 shows a giant Ly- α cloud, which is associated with the radio galaxy 1243+036 at $z = 3.6$ (van Ojik et al. 1996). Several more examples are discussed by Gopal-Krishna et al. (1995) and GW00. A plausible explanation for most such clouds is that the ejection of radio jets, probably triggered by the accretion of dusty material onto a collapsed object (e.g., Lynden-Bell 1996), blows an expanding radio cavity through a pre-existing giant circum-galactic nebula with a two-phase temperature/density structure (Rees 1988). This creates a cocoon of nebular gas around the cavity, which is compressed, heated and accelerated from within by the expanding radio lobes and their bow-shocks. Denser clumps in this ionized cocoon manifest themselves as the Ly- α nebula. (A similar model was proposed by Meisenheimer & Hippelein (1992) to explain the [O II] λ 3727 nebulosity around the radio galaxy 3C368.) The importance of radio lobes in the creation of giant Ly- α nebulae




Figure 2: The $\text{Ly}\alpha$ image of 1243+036 in contours with the 8.3 GHz VLA map in greyscale superimposed. The bending point of the radio structure coincides with enhanced $\text{Ly}\alpha$ emission and a steep rise in the line-width in one of the ‘arms’. Indicated around the edges are the slit positions that were used in the high resolution spectroscopy. Reproduced with permission from van Ojik et al. (1996). ©European Southern Observatory

is further underscored by the finding of Heckman et al. (1991b) who noticed a lack of resolved Ly- α emission among quasars with under-developed radio lobes (i.e., $\leq 10\text{kpc}$). More recently, Giraud et al. (1996) have argued that nuclear UV photons can be an important, and probably dominant, source of excitation for even such giant nebulae (see, also, van Ojik et al. 1996).

In the now standard hierarchical models of the growth of cosmic structure, such Ly- α clouds and newly forming galaxies will be portions of the web of sheets and filaments of baryonic matter collecting within dark matter halos (e.g., Davé et al. 2001).

6.1. Asymmetry of Lyman- α halos as a clue to the sidedness

Discounting the virtual impossibility that in all radio galaxies, the disk is perfectly planar and, moreover, viewed precisely edge-on, one expects to find cases where the Ly- α halo associated with the far-side radio lobe is at least partly hidden behind the disk, and is thus effectively obscured. If, indeed, the Ly- α brightness asymmetry about the nucleus is because the brighter part of the cloud is associated with the *approaching* radio lobe (and hence not obscured by the central dusty disk), then one would expect to detect in the kinematics of the Ly- α cloud a signature of the lobe's motion in our direction.

Such a distinct signature, namely, a negative velocity gradient (blueshift) along the Ly- α cloud from the nucleus toward the outer edge of the lobe, has indeed been found in the case of those high- z RGs which exhibit a distinctly asymmetric Ly- α brightness about the nucleus *and* for which a velocity profile along the cloud is available. These galaxies are: 3C294 ($z = 1.82$, McCarthy et al. 1990), OTL 0852+124 ($z = 2.47$, Gopal-Krishna et al. 1995), 8C 1435+635 ($z = 4.25$, Lacy et al. 1994; Spinrad, Dey & Graham 1995) and 1243+036 ($z = 3.57$, van Ojik et al. 1996). The case of the $z = 3.8$ galaxy 4C41.17 is somewhat confusing, however. The brightest part of the Ly- α cloud which extends to the northeast of the nucleus by $\sim 3''$, appears, instead, to be *redshifted* relative to the nucleus, though the velocity gradient reverses still further to the east (Hippelein & Meisenheimer 1993; Chambers, Miley & van Breugel 1990). Nonetheless, the detection of a radio jet on the eastern side of the nucleus (Carilli, Owen & Harris 1994) strongly supports our case that the brighter Ly- α halo seen on the eastern side is associated with the approaching radio lobe. Further discussion on this point can be found in GW00.

7. Powerful radio sources with one-sided radio structure?

If indeed, going by the standard wisdom, radio lobes have no bulk relativistic motion, the existence of sources with their entire radio structure on just one side of the parent galaxy would furnish strong evidence for a central engine capable of ejecting a one-sided jet (§7.2). As a class, such one-sided radio quasars were first highlighted and christened 'D2 quasars' by Macdonald & Miley (1971). In subsequent deeper radio imaging of the candidates, the 'missing' lobe has been spotted in most cases (e.g., Saikia et al. 1990, and references therein), strengthening the possibility that the D2 quasars simply populate the tail of the lobe flux-ratio distribution. That the extreme flux ratios may actually arise from orientation dependent factors is intimated by the scarcity of such sources among RGs and lobe-dominated radio sources, in general (e.g., LHG; Saikia et al. 1990; Browne & Perley 1986).

7.1. The problem of 3C273

One source which continues to queer the pitch by steadfastly frustrating all attempts to detect any lobe associated with the putative counter-jet is 3C 273, the prototypical D2 quasar. In spite of being imaged with a dynamic range of up to 10,000:1, no convincing detection of a counterpart

to the main radio lobe has been possible (Conway et al. 1981; Davis et al. 1985). Is it then the rare example of truly one-sided jet? Or, alternatively, is in this source even the lobe advancing relativistically and at a small angle to the line-of-sight? Since this may seem unlikely, in view of the rather inconspicuous appearance of the lobe even around the approaching jet, a further alternative has been suggested which obviates the need for a highly relativistic lobe velocity, while still admitting the usual two-sided jet ejection scenario. Basically, in this scenario, 3C 273 is envisioned to be a very energetic, young ($\sim 10^6$ yr) source, with its highly collimated beams, pointed close to the line-of-sight ($\theta \leq 20^\circ$) and advancing rapidly ($\beta \geq 0.6$). Such a rapid advance of the beam is not conducive to lobe formation. Furthermore, due to the transit-time delay, any lobe associated with the tip of the putative counter-jet would be monitored when it was a few times younger than the observed approaching lobe, i.e., at a stage which was probably too early for lobe formation (Kundt & Gopal-Krishna 1981, 1985; Morrison, Roberts & Sadun 1984; Morrison & Sadun 1992; Stawarz 2004). Since this explanation requires several special circumstances, including a close orientation, young age, and a fairly rapid jet advance, the extreme rarity of 3C273-like objects would not be too surprising.

7.2. Producing intrinsically one-sided jets

Although only about one-third of all well mapped radio sources exhibit both jets on the kiloparsec-scale, while another third allow us to view one jet, and the remainder, neither (e.g. Bridle & Perley 1984), the idea that all sources contain essentially symmetrical pairs of jets is inspired by the global symmetry of their radio lobes. The twin-beam picture of Blandford & Rees (1974) suggested that an initially isotropic ejection of hot relativistic plasma from a galactic core could be channelled through de Leval nozzles established in a flattened surrounding medium, into a pair of equally powerful jets. Though the discovery from VLBI of one-sided jets, already on the parsec-scale, showed that this picture was probably not relevant on the ~ 100 pc scale, as envisioned by Blandford & Rees, the possibility that this mechanism played some role on smaller scales could not be excluded (Wiita 1978). Even if the jets are produced in the immediate environs of the assumed supermassive black holes and are intrinsically symmetric on that (innermost) sub-pc scale, interactions with a possibly asymmetric ambient medium on a wide range of scales could easily modify the effective thrusts and/or opening angles of the jets, thus engendering an intrinsic asymmetry.

Wiita & Siah (1981) demonstrated that even a modest offset of the central source from the center of mass of a confining ISM cloud might choke or greatly weaken one of the jets, while allowing the other to burst out. The possible orbiting of the jet emitting engine about the center of the confining cloud could imply that the jets emerged alternately on one side and then on the other, yielding the rough symmetry as a time-averaged effect (e.g., Saikia & Wiita 1982). Icke (1983) suggested a hydrodynamical “clamshell” mechanism that could possibly produce such inversions in the central regions of galaxies. Rudnick & Edgar (1984) reported a significant number of sources with asymmetric radio features that appeared to mirror this “flip-flop” phenomenon (see, also, LHG). The possibility of a binary black hole system in the active nucleus has also been proposed as a way of producing alternating side ejections (Begelman, Blandford & Rees 1980; Komberg 1994). The apparent need for continuous reacceleration of synchrotron emitting electrons because of their short life-times is, however, a strong argument in favor of the vast majority of sources being continually fed on both sides, even if not with a perfect balance.

A discussion of the various mechanisms proposed for the launching and initial collimation of the jets is beyond the scope of this article (see Wiita 1990, 1996 for reviews); here we merely note that essentially all those models are based on accretion disks around supermassive black holes, invoke magnetic fields and tend to produce quite symmetrical, oppositely directed jets. Still, the

possibility that asymmetric jets could be produced by magnetically dominated mechanisms has been discussed (e.g., Bodo et al. 1990; Bisnovatyi-Kogan & Moisenko 1992). The most detailed proposal along these lines is the model of Wang, Sulkanen, & Lovelace (1992), which allows for magnetized disks asymmetric about the mid-plane that can produce very asymmetrical jets. The Wang et al. models are field-dominated and carry energy away from the disk mostly in the Poynting flux of electromagnetic fields.

Other evidence for intrinsically asymmetrical jets is presented by Fraix-Burnet (1992), who points out that Doppler favoritism may not be adequate to explain one-sided jets in sources like M87, NGC 6251 and Cygnus A, whose jets are probably oriented close to the plane of the sky (see, however, Jones 2000). Fraix-Burnet (1992) argues that while a pair of jets is probably always launched, in many cases only one is observed. In stark contrast to the usual assumptions about the consequences of jet interactions with external media (§§5, 7.5), in this scheme the visible jet is supposed to be interacting less with the ISM than is its invisible counterpart. The idea behind this radical interpretation is that if magnetohydrodynamical turbulence produces most of the synchrotron emission from a jet, it may be enhanced if the boundary layer is thinner, i.e., when the jet does not interact intensively with the ISM.

7.3. The apparent asymmetry of jets in powerful radio sources

The asymmetry of optical jets on the kiloparsec-scale was the first striking illustration of asymmetry in RGs (i.e., M87), even preceding the advent of radio astronomy (§1). The initial explanation invoking a bulk relativistic motion of the radiating plasma has received mounting support from observations in different wavebands (Begelman, Blandford & Rees 1984; Scheuer 1987; Urry & Padovani 1995; Cohen & Kellermann 1995; Kellermann et al. 2004). Some of the well known evidences for relativistic motion in both parsec and kiloparsec-scale jets include: the apparent superluminal motion of the parsec-scale jets and their correlated sidedness with the *kiloparsec*-scale jets; the apparent deficit of inverse Compton boosted X-rays from the compact jets; the lobe-depolarization asymmetry of quasars (§3); the intra-day radio variations; the rapid time variability of intense γ -ray emission from blazars; and the correlation of radio structural asymmetry and distortion with the core prominence. Instead of repeating those extensive arguments here, we refer the readers to some review article (e.g. Bridle 1996; Laing 1995; Bicknell 1995; Urry & Padovani 1995; Gopal-Krishna 1995; Chiaberge 2004). Additionally, we recall the growing recognition of the free-free absorption by the nuclear disk as a possible cause of the apparent asymmetry of the parsec-scale radio jets (e.g., Vermeulen, Readhead & Backer 1994; Romney et al. 1995; Matveyenko et al. 1980; 1996).

Here, we briefly recount some observational clues which serve to reveal (indirectly) the presence of collimated energy beams on the kiloparsec-scale, even though the beams are not seen directly. These findings further exemplify that non-detection of jets (a possible cause of jet asymmetry) could well be a mere illusion (however, see the recent detailed optical/Near-IR/UV imaging study of the M87 jet by Meisenheimer, Röser & Schlötelburg 1996; also, van Groningen, Miley, & Norman 1980).

7.4. Invisible beams: Doppler-hidden, or simply non-existent?

Two oft-cited, non-statistical evidences for the existence of ‘invisible’ jets are:

- (i) The discovery of a nonthermal optical synchrotron hot-spot at the expected end-point of the (invisible) counter-jet in M87 (Stiavelli et al. 1992; Sparks et al. 1992). In that the lifetimes of the optically radiating electrons are exceedingly short (≤ 100 yr; Meisenheimer et al. 1996), an active counter-jet has been inferred.
- (ii) From long-slit spectroscopy of the region of the (invisible) counter-jet in the superluminal

galaxy 3C 120, Axon et al. (1989) have found evidence for a ‘cocoon’ of thermal gas which contains signatures of kinematic disturbance arising from the passage of the putative counter-jet through the cocoon.

7.5. Does asymmetric dissipation cause apparent jet asymmetry?

Numerous authors have advocated the possibility that the apparent asymmetry of jets, instead of necessarily arising from relativistic beaming of the approaching jet, could even be caused through an enhanced dissipation of one of the two jets. Conceivably, this could occur in cases where the host galaxy has an asymmetric environment, leading to a more rapid entrainment of the ambient thermal plasma into the relativistic jet on that side (e.g., Bicknell 1984). It is of considerable interest to examine this possibility for the origin of jet asymmetry through jet dissipation, particularly for the case of powerful one-sided jets observed in quasars. The dissipation mechanism is also commonly invoked to explain the absence of hot-spots in low power (FR I) radio galaxies in which the presence of two-sided jets at large distance from the nucleus is interpreted as a manifestation of the jet’s deceleration through the postulated entrainment process (e.g., Bicknell 1994; Komissarov 1994; Laing 1995).

A second possibility proposed for the lack of hot-spots invokes a slowing down of the jet’s front to trans-sonic velocity, leading to its decollimation and the consequent weakening of the Mach disk (e.g., Gopal-Krishna & Wiita 1988; Gopal-Krishna 1991; Roland, Lehoucq & Pelletier 1992; Blandford 1996). Thus, the relativistic jet’s plasma is henceforth discharged into a non-relativistic, plume-like outflow, rather than into a well confined terminal hot-spot. We have shown that this transition may be important in converting FR II sources into FR I’s (Gopal-Krishna & Wiita 2001).

To test these alternatives we (Gopal-Krishna, Wiita, & Hooda 1996) have examined a sample of 8 ‘weak-headed-quasars’ (WHQs) which seem to form a significant fraction ($\sim 10\%$) of a representative sample of highly luminous radio quasars located at high-redshifts ($z \geq 1.5$) and imaged with the VLA at $\lambda = 6$ cm (Lonsdale, Barthel & Miley 1993). The peculiarity marking these 8 sources is that their (one-sided) jets are very prominent but do not terminate in a conspicuous hot-spot. On phenomenological grounds we argued that the observed lack of two-sided jets among the WHQs seems to be more consistent with the second alternative mentioned above (Gopal-Krishna et al. 1996). For, if the entrainment/dissipation model were true, the jet’s visibility would not be determined solely by beaming effects and therefore roughly half of the 8 jets seen should be on the far side of the nucleus and still be visible. But in none of the 8 cases is the putative approaching counterpart of any such misdirected jet detected as well (even though the radiation from any such counterpart would be Doppler boosted toward us). The non-detection of two-sided jets in any of the WHQs supports the Mach-disk decay picture; we argued that various other ways of explaining the result (e.g., hot-spot emission beamed away from us; hot-spot having a much steeper spectral index than the jet; very asymmetric ambient gas distributions) are all quite implausible.

Under the beaming hypothesis, the jetted side is always the one nearer to us, so it is seen at a later stage in the source evolution. This time lag would facilitate the viewing of the shock-front’s disruption and weakening of the hot-spot on the jet’s side. Non-relativistic numerical simulations show that the Mach-disk decay and hot-spot fading can occur, but the more appropriate relativistic simulations have yet to be performed. Additional good quality maps of high- z quasars should yield a larger sample of WHQs, which would permit a more persuasive evaluation of the various competing mechanisms proposed.

8. Conclusions

Over the past three decades astronomers have successfully wielded the radio source asymmetries (or, the lack thereof) as a double-edged knife to attack the twin issues of radio source formation and evolution. It would be too ambitious on our part to attempt a survey of the enormous extant literature on this subject, let alone review it in totality. Therefore we have endeavored here to focus on just a few selected facets of the problem, which we hope serve to highlight the complex interplay between the intrinsic, environmental and viewing-dependent factors, leading to the various manifestations of the asymmetries reported in the literature.

Independent analyses/interpretations of the observed asymmetries in the lengths, luminosities, spectra, spectral curvature, depolarization, optical/UV surface brightness and non-colinearity of the twin-lobes, have provided a number of key leads and several important conclusions, especially when the relativistic speed of the jets is taken into account. As discussed in the previous sections, these findings form a crucial part of the growing body of evidence in favor of the orientation-based unification theories. The outcome of the asymmetry studies now also needs to be harnessed for exploring the evolutionary connections between the two main morphological classes of radio galaxies (FR I and FR II), and for probing the relationships between the dimensional sequence represented by compact-symmetric objects (CSOs), Gigahertz-peaked-spectrum (GPS) sources, compact-steep-spectrum (CSS) sources, intermediate-symmetric-objects (ISO), classical RGs, and giant radio galaxies (GRG) (e.g., Snellen et al. 2000).

Investigations of radio source asymmetries have raised several outstanding questions whose resolution will require continued confrontation of thought with observations. We sum up this review with the proverbial set of 10 posers related to this topic, though the careful reader may note that we have squeezed in several extra follow-up questions:

- Are the mild hot-spot speeds ($\approx c/100$), as indicated by the Q -distributions of radio-luminous quasars, consistent with the markedly stronger asymmetry observed for quasars (compared to RGs), if orientation is the only difference between the two?
- Do powerful radio sources at high redshifts have more distorted radio structures, compared to their less distant counterparts?
- Typically, how large is the ‘backflow’ speed in classical double radio sources? What is its dependence on the speed of the hot-spot?
- For FR II double radio sources, how reliably can the apparent asymmetry of the Ly- α halo be used to identify the near-side lobe? How does the Ly- α asymmetry correlate with the lobe depolarization asymmetry (which is a fairly robust indicator of the sidedness)?
- How different are the spectral-index asymmetries for the radio lobes and the hot-spots? Such information may provide vital clues for disentangling intrinsic, environmental and orientational effects on the apparent spectral-index asymmetry.
- How significant is the contribution of the parent galaxy’s motion to the lobe-length asymmetry?
- Will the correlated radio-optical asymmetry turn out to be a weaker effect if red emission-line images of the EELR are used for quantifying the optical asymmetry? Such a finding would support the notion that the optical asymmetry is primarily a result of asymmetric dust obscuration (rather than due to an asymmetric line-emission).
- How genuine is the ‘mystique’ of 3C273? Can nature make truly one-sided jets?
- Are there RGs (located in extremely rarefied environments) whose jets are still boring their way at relativistic speed on the kiloparsec scale (and consequently, have not yet formed their hot-spots and lobes)?
- What is the late phase of the evolutionary scenario for jets? What makes their hot-spots finally decay (e.g., weak-headed-quasars: WHQs)? Does their decay finally destabilize the jets themselves?

This question also has a strong bearing on the issue of unification of the two Fanaroff-Riley morphological classes.

Acknowledgment: We offer our apologies to all those authors whose work was not cited in this article, in spite of its relevance; these omissions are not merely due to our ignorance, but also arose from the combination of limited space and the vastness of the literature on this topic. We thank J.P. Leahy and H. Röttgering for providing the electronic versions of Figs. 1 and 2, respectively. This work was supported in part by NASA grant NAG5-3098 and Research Program Enhancement funds to the Program in Extragalactic Astronomy at Georgia State University. PJW is grateful for continuing hospitality at the Department of Astrophysical Sciences at Princeton University.

References

- Antonucci, R., & Barvainis, R. 1990, *ApJ*, 363, L17
- Arshakian, T.G., & Longair, M.S. 2000, *MNRAS*, 311, 846
- Axon, D.J., Unger, S.W., Pedlar, A., Meurs, E.J.A., Whittle, D.M., & Ward, M.J. 1989, *Nature*, 341, 631
- Baade, W. 1956, *ApJ*, 123, 550
- Banhatti, D.G. 1980, *A&A*, 84, 112
- Barger, A.J., et al. 2001, *AJ*, 122, 2177
- Barthel, P.D. 1989, *ApJ*, 336, 606
- Barthel, P.D., & Miley, G.K. 1988, *Nature*, 333, 319
- Baryshev, Yu., & Teerikorpi, P. 1995, *A&A*, 295, 11
- Begelman, M.C., Blandford, R.D., & Rees, M.J. 1980, *Nature*, 287, 307
- Begelman, M.C., Blandford, R.D., & Rees, M.J. 1984, *Rev. Mod. Phys.* 56, 255
- Best, P.N., Bailer, D.M., Longair, M.S., & Riley, J.M. 1995, *MNRAS*, 275, 1171
- Bicknell, G. 1984, *ApJ*, 286, 68
- Bicknell, G. 1994, *ApJ*, 422, 542
- Bicknell, G. 1995, *ApJS*, 101, 29
- Bisnovatyi-Kogan, G.S. & Moisenko, S.G. 1992, *Astron. Zh.*, 36, 285
- Black, A.R.S., Baum, S.A., Leahy, J.P., Perley, R.A., Riley, J.M., & Scheuer, P.A.G. 1992, *MNRAS*, 256, 186
- Blandford, R.D. 1996, in *Cygnus A – Study of a Radio Galaxy*, C. Carilli & D. Harris, Cambridge: Cambridge Univ. Press, 264
- Blandford, R.D., & Rees, M.J. 1974, *MNRAS*, 169, 395
- Blundell, K.M., & Alexander, P. 1994, *MNRAS*, 267, 241
- Blundell, K.M., & Rawlings, S. 2000, *AJ*, 119, 1111
- Blundell, K.M., & Rawlings, S. 2001, in *Particles and Fields in Radio Galaxies*, ASP Conf. Ser. 250, R.A. Laing & K.M. Blundell, San Francisco: ASP, 363
- Blundell, K.M., Rawlings, S., & Willott, C.J. 1999, *AJ*, 117, 677
- Bodo, G., Chagelishvili, G.D., Ferrari, A., Lominadze, J.G., & Trussoni, E. 1990, in *Plasma Astrophysics*, ESA SP-311, T.D. Guyenne & J.J. Hune, Paris: ESA, 273
- Bregman, J.N., Hogg, D.E., & Roberts, M.S. 1995, *ApJ*, 441, 561
- Bridle, A.H. 1996, in *Energy Transport in Radio Galaxies and Quasars*, ASP Conf. Ser. 100, P.E. Hardee, A.H. Bridle, & J.A. Zensus, San Francisco: ASP, 383
- Bridle, A.H., & Perley, R.A. 1984, *ARA&A*, 22, 319
- Bridle, A.H., Laing, R.A., Scheuer, P.A.G. & Turner, S. 1994, in *The Physics of Active Galaxies*, G.V. Bicknell, M. Dopita, P. Quinn, ASP Conf. Ser. 54, San Francisco: ASP, 187
- Browne, I.W.A., & Perley, R.A. 1986, *MNRAS*, 222, 149
- Capetti, A., Morganti, R., Parma, P., & Fanti, R. 1993, *A&AS*, 99, 407

- Chambers, K.G., Miley, G.K., van Breugel, W.J.M. 1990, *ApJ*, 363, 21
- Carilli, C. L., Owen, F. N. & Harris, D. E. 1994, *AJ*, 107, 480
- Cen, R. & Ostriker, J. 1999, *ApJ*, 519, L109
- Chiaberge, M. 2004, *astro-ph/0405255*
- Chini, R., & Krügel, E. 1994, *A&A*, 288, L33
- Chyży, K.T., & Zięba, S. 1995, *A&A*, 303, 420
- Cimatti, A., di Serego Alighieri, S., Fosbury, R.A.E., Salvati, M., & Taylor, D. 1993, *MNRAS*, 264, 421
- Clarke, D.A. 1996, in *Energy Transport in Radio Galaxies and Quasars*, ASP Conf. Ser. 100, P.E. Hardee, A.H. Bridle, & J.A. Zensus, San Francisco: ASP, 311
- Cohen, M.H. & Kellermann, K.I. 1995, *Proc. Natl. Acad. Sci. USA*, 92, 11339
- Colina, L., & de Juan, L. 1995, *ApJ*, 448, 548
- Conway, R.G., Davis, R.J., Foley, A.R., & Ray, T.P. 1981, *Nature*, 294, 540
- Cowie, L.L., Hu, E.M., & Songaila, A. 1996, *Nature*, 377, 603
- Curtis, H.D. 1918, *Lick Obs. Publ.*, 13, 11
- Davé, R., et al. 2001, *ApJ*, 355, 416
- Davis, R.J., Muxlow, T.W.B., & Conway, R.G. 1985, *Nature*, 318, 343
- Dennett-Thorpe, J., Bridle, A.H., Scheuer, P.A.G. Laing, R.A., Leahy, J.P. 1997, *MNRAS*, 289, 753
- De Young, D.S., & Axford, W.I. 1967, *Nature*, 216, 129
- Dunlop, J.S., Hughes, D.H., Robbins, S., Eales, S.A., & Ward, M.J. 1994, *Nature*, 370, 347
- Eilek, J.A., & Arendt, P.N. 1996, *ApJ*, 457, 150
- Ensmann, L.M., & Ulvestad, J.S. 1984, *AJ*, 89, 1275
- Evrard, A.E., Summers, F.J., & Davis, M. 1994, *ApJ*, 422, 11
- Fokker, A.D. 1986, *A&A*, 156, 315
- Fomalont, E.B. 1969, *ApJ*, 157, 1027
- Fraix-Burnet, D. 1992, *A&A*, 259, 445
- Fukugita, M., Hogan, C.J., & Peebles, P.J.E. 1996, *Nature*, 381, 489
- Garrington S.T., & Conway, R.G. 1991, *MNRAS* 259, 198
- Garrington S.T., Conway, R.G., & Leahy, J.P. 1991, *MNRAS*, 250, 171
- Garrington, S.T., Leahy, J.P., Conway, R.G., & Laing, R.A. 1988, *Nature*, 331, 147
- Georganopoulos, M., & Kazanas, D. 2003, *ApJ*, 589, L5
- Giraud, E., Melnick, J., Gopal-Krishna, Oliveira, C. & Kulkarni, V.K. 1996, *A&A*, 309, 733
- Gopal-Krishna 1980, *A&A*, 86, L1
- Gopal-Krishna 1991, *A&A*, 248, 415
- Gopal-Krishna 1995, *Proc. Nat. Acad. Sci. USA*, 92, 11399
- Gopal-Krishna, Giraud, E., Melnick, J., & Della Valle, M. 1995, *A&A*, 303, 705
- Gopal-Krishna, Kulkarni, V.K., & Wiita, P.J. 1996, *ApJ*, 463, L1
- Gopal-Krishna & Nath, B. 1997, *A&A*, 326, 45
- Gopal-Krishna & Wiita, P.J. 1988, *Nature*, 333, 49
- Gopal-Krishna & Wiita, P.J. 1996, *ApJ*, 467, 191 (GW96)
- Gopal-Krishna & Wiita, P.J. 2000, *ApJ*, 529, 189 (GW00)
- Gopal-Krishna & Wiita, P.J. 2001, *A&A*, 373, 100
- Gopal-Krishna, Wiita, P.J., & Hooda, J.S. 1996, *A&A*, 316, L13
- Gopal-Krishna, Wiita, P.J., & Saripalli, L. 1989, *MNRAS*, 239, 173
- Goudfrooij, P. & de Jong, T. 1995, *A&A*, 298, 784
- Hargrave, P.J., & McEllin, M. 1975, *MNRAS*, 173, 37
- Hargrave, P.J., & Ryle, M. 1974, *MNRAS*, 166, 305
- Heckman, T.M., Lehnert, M.D., van Breugel, W., & Miley, G.K. 1991a, *ApJ*, 370, 78
- Heckman, T.M., Lehnert, M.D., Miley, G.K., & van Breugel, W. 1991b, *ApJ*, 381, 373

- Hintzen, P., & Scott, J.S. 1978, ApJ, 224, L47
- Hippelein, H. & Meisenheimer, K. 1993, Nature, 362, 224
- Hooda, J.S., & Wiita, P.J. 1998, ApJ, 493, 81
- Hough, D.H. & Readhead, A.C.S. 1989, AJ, 98, 1208
- Icke, V. 1983, ApJ, 265, 648
- Ingham, W., & Morrison, P. 1975, MNRAS, 173, 569
- Ishwara-Chandra, C.H., & Saikia, D.J. 2002, New Astr. Rev., 46, 71
- Jennison, R.C., & Das Gupta, M.K. 1953, Nature, 172, 996
- Jeyakumar, S., Wiita, P.J., Saikia, D.J., & Hooda, J.S., 2004, A&A, in press
- Johnson, R.A., Leahy, J.P., & Garrington, S.T. 1995, MNRAS, 273, 877
- Kapahi, V.K. 1989, AJ, 97, 1
- Kapahi, V.K. 1990a, in Parsec-scale Radio Jets, J.A. Zensus & T.J. Pearson, Cambridge: Cambridge University Press, 304
- Kapahi, V.K. 1990b, Current Sci., 59, 561
- Kapahi, V.K., & Saikia, D.J. 1982, J. Ap. Astr., 3, 465
- Kellermann, K.I. et al. 2004, ApJ, 609, 539
- Komberg, B.V. 1994, Astr. Rep., 38, 617
- Kommissarov, S.S. 1994, MNRAS, 269, 394
- Kotanyi, C.G., & Ekers, R.D. 1979, A&A, 73, L1
- Kundt, W., & Gopal-Krishna, 1980, Nature, 288, 149
- Kundt, W., & Gopal-Krishna, 1981, Ap. Space Sci., 75, 257
- Kundt, W., & Gopal-Krishna, 1986, J. Ap. Astr., 7, 225
- Lacy, M., et al. 1994, MNRAS, 271, 504
- Laing, R.A., 1988, Nature, 331, 149
- Laing, R.A. 1995, Proc. Natl. Acad. Sci. USA, 92, 11413
- Lawrence, A. 1991, MNRAS, 252, 586
- Leahy, J.P., Muxlow, T.W.B., & Stephens, P.W. 1989, MNRAS, 239, 401
- Leahy, J.P., & Perley, R.A. 1991, AJ, 102, 537
- Lister, M.L., Gower, A.C., & Hutchings, J.B. 1994, AJ, 108, 821
- Lister, M.L., Hutchings, J.B., & Gower, A.C. 1994, ApJ, 427, 125 (LHG)
- Liu, R. & Pooley G.G. 1991a, MNRAS, 249, 343
- Liu, R. & Pooley G.G. 1991b, MNRAS, 253, 669
- Liu, R., Pooley, G.G., & Riley, J.M. 1992, MNRAS, 257, 545
- Longair, M.S., & Riley, J.M. 1979, MNRAS, 188, 625
- Longair, M.S., Ryle, M., & Scheuer, P.A.G. 1973, MNRAS, 164, 243
- Lonsdale, C.J., Barthel, P.D., & Miley, G.K. 1993, ApJS, 87, 63
- Lynden-Bell, D. 1996, MNRAS, 279, 389
- Macdonald, G.H., & Miley, G.K. 1971, ApJ, 164, 237
- Mackay, C.D. 1971, MNRAS, 154, 209
- Mackay, C.D. 1973, MNRAS, 162, 1
- Macklin, J.T. 1981, MNRAS, 196, 967
- Mangalam, A.V. & Gopal-Krishna 1995, MNRAS 275, 979
- Marconi, A., Risaliti, G., Gilli, R., Hunt, L.K., Maiolino, R., Salvati, M. 2004, MNRAS, 351, 169
- Matveyenko, L.I. et al. 1980, SvAL, 6, 42
- Matveyenko, L.I., Pauliny-Toth, I.I.K., Graham, D.A., Sherwood, W.A., B  ath, L.B., & Kus, A.D. 1996, Astr. Lett., 22, 14
- McCarthy, P.J., Spinrad, H., van Breugel, W., Liebert, J., Dickinson, M., Djorgovski, & Eisenhardt, P. 1990, ApJ, 365, 487

- McCarthy, P.J. & van Breugel, W. 1989, in *Extranuclear Activity in Galaxies*, E.J.A. Meurs & R.A.E. Fosbury, Garching: ESO, 55
- McCarthy, P.J., van Breugel, W., & Kapahi, V.K. 1991, *ApJ*, 371, 478 (MvBK)
- Meisenheimer, K. & Hippelein, H. 1992, *A&A*, 264, 465
- Meisenheimer, K., Röser, H.-J. & Schlötelburg, M. 1996, *A&A*, 307, 61
- Mellema, G., Kurk, J.D., & Röttgering, H.J.A. 2002, *A&A*, 395, L13
- Miley, G.K. 1980, *ARA&A*, 18, 165
- Miller, C.J., Nichol, R.C., Gómez, P.L., Hopkins, A.M., & Bernardi, M. 2003, *ApJ*, 597, 142
- Morisawa, K. & Takahara, F. 1987, *MNRAS*, 228, 745
- Morrison, P., Roberts, D., & Sadun, A. 1984, *ApJ*, 280, 483
- Morrison, P., & Sadun, A. 1992, *MNRAS*, 254, 488
- Norman, M.L. 1996, in *Energy Transport in Radio Galaxies and Quasars*, ASP Conf. Ser. 100, P.E. Hardee, A.H. Bridle, & J.A. Zensus, San Francisco: ASP, 319
- Parma, P., de Ruiter, H.R., Mack, K.-H., van Breugel, W., Dey, A., Fanti, R., & Klein, U. 1994, *A&A*, 311, 49
- Peacock, J.A. 1987, in *Astrophysical Jets and their Engines*, W. Kundt, Dordrecht: Reidel, 185
- Pedetty, J.A., Rudnick, L., McCarthy, P.J., & Spinrad, H. 1989, *AJ*, 97, 647
- Rees, M.J. 1988, *MNRAS*, 239, 1P
- Rees, M.J., Begelman, M. C., Blandford, R. D. & Phinney, E. S. 1982, *Nature*, 295, 17
- Riley, J.M & Jenkins, C.J. 1977, in *Radio Astronomy and Cosmology*, IAU Symp. 74, D.L. Jauncey, Dordrecht: Reidel, 237
- Roland, J., Lehoucq, R., & Pelletier, G. 1992, in *Extragalactic Radio Sources—from Beams to Jets*, J. Roland, H. Sol, G. Pelletier, Cambridge: Cambridge University Press, 294
- Romney, J.D., Biretta, J.M., Dhawan, V., Kellermann, K.I., Vermeulen, R.C., & Walker, R.C. 1995, *Proc. Natl. Acad. Sci. USA*, 92, 11360
- Rudnick, L., & Edgar, B.K. 1984, *ApJ*, 279, 74
- Rudnick, L., Katz-Stone, D.M., & Anderson, M.C. 1994, *ApJS*, 90, 955
- Ryle, M., & Longair, M.S. 1967, *MNRAS*, 136, 123
- Ryś, S. 1994, *A&A*, 281, 15
- Saikia, D.J. 1984, *MNRAS*, 245, 408
- Saikia, D.J., Junor, W., Cornwell, T.J., Muxlow, T.W.B., & Shastri, P. 1990, *MNRAS*, 245, 408
- Saikia, D.J., Jeyakumar, S.K., Wiita, P.J., Sanghera, H.S., & Spencer, R.E. 1995, *MNRAS*, 276, 1215
- Saikia, D.J., Jeyakumar, S.K., Mantovani, F., Salter, C.J., Spencer, R.E., Thomasson, P., & Wiita, P.J. 2003, *PASA*, 20, 50
- Saikia, D.J., & Wiita, P.J. 1982, *MNRAS*, 200, 83
- Saripalli, L., Gopal-Krishna, Reich, W., & Kühr, H. 1986, *A&A*, 170, 20
- Scheuer, P.A.G. 1974, *MNRAS*, 166, 513
- Scheuer, P.A.G. 1982, in *Extragalactic Radio Sources*, IAU Symp. 97, D.S. Heeschen, C.M. Wade, Dordrecht: Reidel, 163
- Scheuer, P.A.G. 1987, in *Astrophysical Jets and their Engines*, W. Kundt, Dordrecht: Reidel, 129
- Scheuer, P.A.G. 1995, *MNRAS*, 277, 331
- Siah, J., & Wiita, P.J. 1990, *ApJ*, 363, 411
- Singal, A.K. 1988, *MNRAS*, 233, 87
- Slysh, V.I. 1966, *Sov. Astr.*, 9, 533
- Snellen, I.A.G., Schilizzi, R.T., Miley, G.K., de Bruyn, A.G., Bremer, M.N. & Röttgering, H.J.A. 2000, *MNRAS*, 319, 445
- Sofue, Y., & Wakamatsu, K. 1992, *PASJ*, 44, L23
- Sparks, W.B., Fraix-Burnet, D., Macchetto, F., & Owen, F.N. 1992, *Nature*, 355, 804
- Spinrad, H., Dey, A., & Graham, J.R. 1995, *ApJ*, 438, L51

- Stawarz, L. 2004, *ApJ*, 613, 119
- Stiaveli, M., Biretta, J., Moller, P., Zeilinger, W.W. 1992, *Nature*, 355, 802
- Strom, R.G. & Jägers, G. 1988, *A&A*, 194, 79
- Swarup, G., & Banhatti, D.G. 1981, *MNRAS*, 194, 1025
- Teerikorpi, P. 1984, *A&A*, 132, 179
- Teerikorpi, P. 1986, *A&A*, 164, L11
- Tribble, P.C. 1992, *MNRAS*, 256, 281
- Urry, C.M., & Padovani, P. 1995, *PASP*, 107, 803
- Valtonen, M.J. 1979, *ApJ*, 231, 312
- van Groningen, E., Miley, G.K., & Norman, C. 1980, *A&A*, 90, L7
- van Ojik, R., Röttgering, H.J.A., Carilli, C.L., Miley, G.K., Bremer, M.N. & Macchetto, F. 1996, *A&A*, 313, 25
- Vermeulen, R.C., Readhead, A.C.S., & Backer, D.C. 1994, *ApJ*, 429, L41
- Wang, J.C.L., Sulkanen, M.E., & Lovelace, R.V.E. 1992, *ApJ*, 390, 46
- Wang, Z., Wiita, P.J., & Hooda, J.S. 2000, *ApJ*, 534, 201
- Wiita, P.J. 1978, *ApJ*, 221, 41
- Wiita, P.J. 1990, in *Beams and Jets in Astrophysics*, P.A. Hughes, Cambridge: Cambridge University Press, 379
- Wiita, P.J. 1996, in *Energy Transport in Radio Galaxies and Quasars*, ASP Conf. Ser. 100, P.E. Hardee, A.H. Bridle, & J.A. Zensus, San Francisco: ASP, 395
- Wiita, P.J., & Gopal-Krishna 1990, *ApJ*, 353, 476
- Wiita, P.J., & Siah, M.J. 1981, *ApJ*, 243, 710
- Willis, A.G., Wilson, A.S., & Strom, A.G. 1978, *A&A*, 66, L1
- Wills, B.J., Netzer, H., Brotherton, M.S., Han, M., Wills, D., Baldwin, J.A., Ferland, G.J., & Browne, I.W.A. 1993, *ApJ*, 410, 534
- Wilson, A.S. & Colbert, E.J.M. 1995, *ApJ*, 438, 62
- Yee, H.K.C., & Ellingson, E. 1993, *ApJ*, 411, 43
- York, D. G. et al. 2000, *AJ*, 120, 1579
- Zięba, S. & Chyży, K.T. 1991, *A&A*, 241, 22